

Quasiperiodic fields and Bose-Einstein condensation

P. F. Borges,^{*}H. Boschi-Filho[†]and C. Farina[‡]

Instituto de Física, Universidade Federal do Rio de Janeiro

Cidade Universitária, Ilha do Fundão, Caixa Postal 68528

21945-970 Rio de Janeiro, BRAZIL

Abstract

We construct a partition function for fields obeying a quasiperiodic boundary condition at finite temperature, $\psi(0; \vec{x}) = e^{i\theta} \psi(\beta; \vec{x})$, which interpolate continuously that ones corresponding to bosons and fermions and discuss the possibility of condensation for these fields.

PACS: 11.10.Wx; 05.30.-d; 03.65.Db.

Keywords: Bose-Einstein condensation; partition functions.

^{*}e-mail: pborges@if.ufrj.br;

[†]e-mail: boschi@ctp.mit.edu;

[‡]e-mail: farina@if.ufrj.br.

The concept of condensation was proposed by Einstein [1], after the fundamental work of Bose [2] on the statistics of integer spin particles, and its first realization was given by London in a model for the Helium superfluidity [?]. More recently, the interest for this phenomenon has increased because of the experimental results on atomic traps and the direct observation of Bose-Einstein condensation (BEC) on gases [4] signaling for a large number of applications.

Long ago, it was shown for massive bosons in the absence of external fields that BEC can only occur if $N \geq 3$, where N is the space dimension. For massless bosons this condition reduces to $N \geq 2$, although in this case the critical temperature is infinite. These results were first shown in a nonrelativistic context [5] and later generalized the relativistic case [6-8]. Very recently, the extension of these results to curved space-times and the inclusion of non-uniform magnetic fields in the relativistic context were also done [9, 10].

Here, we present a relativistic quantum field theory approach to BEC from which we obtain nonrelativistic results as a particular limit. We start from a partition function identified with a determinant for fields which satisfy a quasiperiodic boundary condition, *i.e.*, a condition which interpolates continuously the periodic (bosonic) and antiperiodic (fermionic) cases at finite temperature. This kind of field is usually considered in lower dimensional systems as in the anyon theory in 2+1 dimensions [11, 12]. Recently, we have shown [13] that the present construction leads to a partition function which is equivalent to the usual ones for anyons, at least for the second virial coefficient [14]. Further within our approach, we also obtain a general condensation condition for fields which interpolate continuously between bosons and fermions and discuss the role of the dimension where the condensation occurs.

Let us start by writing the partition function for a general system described by a Hamiltonian H with chemical potential μ as

$$\mathcal{Z} \equiv \exp\{-\beta\Omega\} = \text{tr } e^{-\beta(H+\mathcal{N}\mu)}, \quad (1)$$

where Ω is the free energy and $\mathcal{N}$ is the conserved charge (conserved quantum number) of the system. As it is well known, for relativistic charged massive bosonic fields one can

express this partition function as a determinant [8, 15], namely,

$$\mathcal{Z}_{Bosons} = \left[\det(-D^2 + M^2) \Big|_P \right]^{-1}, \quad (2)$$

where D^2 is the square of the covariant derivative, including the chemical potential, $D_\nu = (\partial_0 + i\mu, \partial_i)$ and M is the mass of the field. This prescription for the inclusion of the chemical potential μ coincides with the one given by eq. (1) except for the introduction of an imaginary part to $\mathcal{Z}$, which is not physically relevant. So unless otherwise specified, we are considering that the real part of $\mathcal{Z}$ has been taken. Note also that, the chemical potential appears squared in the Klein-Gordon (KG) operator, $-D^2 + M^2$. This is a consequence of representing the partition function in phase space and then integrating over momenta (see [8, 16] for details). The label P means that the eigenvalues of this operator are subjected to periodic boundary conditions and hence they are given by

$$\lambda_{nk} = (\omega_n + i\mu)^2 + \vec{k}^2 + M^2, \quad (3)$$

where $\omega_n = 2n\pi/\beta$, with $n \in \mathbb{Z}$, are the Matsubara frequencies [17] for bosonic fields and $\vec{k} \in \mathbb{R}^N$.

Then, we recall the partition function for a free relativistic fermionic gas. As for the case of bosons, one can write this partition function as a determinant [8, 16]

$$\begin{aligned} \mathcal{Z}_{Fermions} &= \det_D(i\mathcal{D} - M) \Big|_A \\ &= \left[\det(-D^2 + M^2) \Big|_A \right]^{+1}, \end{aligned} \quad (4)$$

where $\det_D$ means the calculation over Dirac indices and the subscript A means that the eigenvalues of each operator are computed with antiperiodic boundary conditions. Hence, for the operator $-D^2 + M^2$, the eigenvalues are given by (3), but now the Matsubara frequencies are $\omega_n = (2\pi/\beta)(n + 1/2)$ with $n \in \mathbb{Z}$.

Now, with the aid of a convenient choice of parameters [13] we are going to use an approach which describe both the bosonic and fermionic partition functions in an unified way. Consider then the generalized determinant

$$\mathcal{Z} = \left[\det(-D^2 + M^2) \Big|_\theta \right]^\sigma, \quad (5)$$

where the subscript θ means that the quasiperiodic finite temperature condition, i.e.,

$$\psi(0; \vec{x}) = e^{i\theta} \psi(\beta; \vec{x}), \quad (6)$$

is assumed and we introduced a new parameter σ to be able to reproduce correctly the bosonic and fermionic particular cases. Observe that when $\theta = 0$ and $\sigma = -1$ we obtain the bosonic partition function, while for $\theta = \pi$ and $\sigma = +1$ we have the fermionic one. The eigenvalues in this case are

$$\lambda_{nk\theta} = \left[\frac{2n\pi}{\beta} + i\mu + \frac{\theta}{\beta} \right]^2 + \vec{k}^2 + M^2. \quad (7)$$

The determinant (5) is a generalization of its quantum mechanical ($0 + 1$) dimensional version (with $\mu = 0$) which has been calculated using Green functions [18] and the zeta function method [19].

Notice that the interpolating parameter θ plays the role of the zeroth component of a topological (temperature-dependent) gauge field $A_\mu = (A_0(\beta), \vec{0})$, since the θ -boundary condition implied that $\omega_n \rightarrow \omega_n + \theta/\beta$, which can be viewed as a shift in the time derivative operator, $\partial_0 \rightarrow \partial_0 + i\theta/\beta$. The discussion of the bosonic (2) and fermionic (4) determinants with a constant topological gauge field, but without relating it to boundary conditions or interpolating statistics, has been presented before [16]. It is important to mention that the parameter θ or equivalently the $A_0(\beta)$ field can not be gauged away because of the nontrivial boundary conditions introduced by the time compactification to the interval $[0, \beta]$. An interesting point that has not been observed before is that the coupling between bosons with the topological field $A_0(\beta)$ suppresses the Bose-Einstein condensation, unless for very specific values of θ , as we are going to show.

It can be shown that the free energy corresponding to the partition function (5), is given by [13]:

$$\begin{aligned} \Omega(\beta, \mu) &\equiv -\frac{1}{\beta} \ln \mathcal{Z} = -\frac{\sigma}{\beta} \ln \det \mathcal{A} \\ &= 4\sigma V \left(\frac{M}{2\pi\beta} \right)^{\frac{N+1}{2}} \sum_{n=1}^{+\infty} \cos(n\theta) \cosh(n\beta\mu) \left(\frac{1}{n} \right)^{\frac{N+1}{2}} K_{\frac{1}{2}(N+1)}(n\beta M), \end{aligned} \quad (8)$$

where $K_\nu(x)$ is the modified Bessel function of the second kind. Note that in the above formula, only the real part of the free energy $\Omega(\beta, \mu)$ is expressed, according to the prescription of introducing the chemical potential as an imaginary time-component gauge potential [8]. If we particularize the parameters σ and θ to the bosonic ($\sigma = -1$ and $\theta = 0$) and fermionic ($\sigma = +1$ and $\theta = \pi$) cases we shall find precisely the results known

in the literature [7, 20, 16]. In a recent work [13], we have shown that this partition function with arbitrary θ leads, in the nonrelativistic limit, to a second virial coefficient which is equivalent to the well known result of Arovas *et. al* [14]. We have also calculated higher order virial coefficients showing good agreement with anyon perturbation theory and numerical calculations [21]. There, we have shown that, for statistical consistency we have to impose that the parameters θ and σ are related by

$$\sigma \equiv \sigma(\theta) = -\cos^{-1} \theta. \quad (9)$$

To see explicitly when the condensation occurs, let us rewrite the free energy (8) as [16]

$$\begin{aligned} \Omega(\beta, \mu) &= -\frac{TV\sigma(\theta)}{(2\pi)^N} \int d^N k \, \Re \left\{ \ln \left[1 - e^{i\theta} e^{-\beta(\omega-\mu)} \right] + (\mu \rightarrow -\mu) \right\} \\ &= -\frac{TV\sigma(\theta)}{2(2\pi)^N} \int d^N k \left\{ \ln \left[1 + e^{-2\beta(\omega-\mu)} - 2 \cos \theta e^{-\beta(\omega-\mu)} \right] + (\mu \rightarrow -\mu) \right\} \end{aligned} \quad (10)$$

where $\omega = \sqrt{\vec{k}^2 + M^2}$. Further, the charge density is given by:

$$\rho \equiv \frac{1}{\beta V} \frac{\partial}{\partial \mu} \ln \mathcal{Z} = -\frac{1}{V} \left(\frac{\partial \Omega}{\partial \mu} \right)_{V,T}. \quad (11)$$

Remind that it goes to infinity when particles condense. From this charge density we can write the general θ -dependent distribution function

$$\begin{aligned} f_\theta(\beta, \pm\mu) &= \sigma(\theta) \Re \left[\frac{1}{1 - e^{-i\theta} e^{\beta(\omega \mp \mu)}} \right] \\ &= -\frac{1}{2 \cos \theta} \left[\frac{\cos \theta - e^{-\beta(\omega \mp \mu)}}{\cosh \beta(\omega \mp \mu) - \cos \theta} \right]. \end{aligned} \quad (12)$$

The condensation condition comes from the divergence in the free energy or equivalently in the corresponding charge density which is also evident in the above distribution function. As the condensation is related to the zero momenta ($\vec{k} = \vec{0}$) state, this implies that $\omega \rightarrow M$ and in the general case, with arbitrary θ we have that this condition is given by

$$1 + e^{-2\beta(M-\mu)} - 2 \cos \theta e^{-\beta(M-\mu)} = 0, \quad (13)$$

or simply

$$\cosh \beta(\mu - M) = \cos \theta, \quad (14)$$

which can only be satisfied for finite temperatures if $\theta = 2n\pi$ (with n integer) and $\mu = M$, simultaneously. This means that the condensation condition in the general case is exactly the same as in the usual BEC.

Then, we conclude that equation (10), or equivalently (8), is well defined¹ for every value of μ , except when $\theta = 2n\pi$ for which $\mu = M$ gives the well known condensation condition. So, we can infer that no condensation occurs for any $\theta \neq 2n\pi$ (n integer). However, this does not forbid the condensation of fermions or θ -fields.

Consider, for instance, that the free θ -field obeying the Klein-Gordon equation is now interacting with another topological field A'_0 , different from the A_0 introduced above to transmutate the statistics of the original bosonic or fermionic field. In this case, the parameter defining the statistics and condensation condition for the θ -field together with the topological interaction can be written as $\Theta = \theta + \theta'$, where θ' is associated with the A'_0 field. Then, the statistical properties in this situation are determined by the parameter Θ . This means that apart from the usual BEC, fermions coupled to topological fields with $\theta' = n\pi$ so that the total phase is $\Theta = 2n\pi$ may possibly condense. Analogously, any θ -field coupled to a topological field with $\theta' = \theta - 2n\pi$, such that the total phase is $\Theta = 2n\pi$ may also condense.

To obtain the critical temperatures, densities and dimensions for the condensate from the above discussion we go back to the free energy given by Eq. (8). Note that this equation contains all the relevant information about our system. However, we are mainly interested in the ultra and nonrelativistic limiting cases.

Let us first consider the ultrarelativistic case. In this situation where the energies involved are much higher than the mass scale (the mass M of the particles) we should take the limit $\beta M \ll 1$, which is easily recognized as a high temperature limit. In this regime we can use the following property of the Bessel functions

$$\lim_{x \rightarrow 0} K_\nu(x) = \Gamma(\nu) 2^{\nu-1} x^{-\nu} ; \quad (\nu > 0) \quad (15)$$

¹Note that from (10) or (12) one can see that there is another possibility of infinite charge density which comes from the choice $\theta = (2n+1)\pi/2$, since $\sigma \rightarrow \pm\infty$, because of relation (9). However, this divergence does not depend on the other physical parameters of the system and then this singularity opposed to the BEC case, seems to be unphysical.

so that the free energy (8) reduces to ($\theta = 2n\pi$, $\sigma(\theta) = -1$)

$$\Omega(\beta, \mu) = -\frac{2V}{(\beta\sqrt{\pi})^{N+1}} \Gamma\left(\frac{N+1}{2}\right) \sum_{n=1}^{+\infty} \cosh(n\beta\mu) \left(\frac{1}{n}\right)^{N+1}. \quad (16)$$

Using now Eq. (11) we find that

$$\rho(\beta, \mu) = \frac{2}{(\sqrt{\pi})^{N+1}} \left(\frac{1}{\beta}\right)^N \Gamma\left(\frac{N+1}{2}\right) \sum_{n=1}^{+\infty} \sinh(n\beta\mu) \left(\frac{1}{n}\right)^N. \quad (17)$$

As we are dealing with the high temperature limit, $\beta M \ll 1$, the condensation condition $\mu \rightarrow M$ also implies $\beta\mu \ll 1$, so that

$$\rho = \frac{2\mu}{\pi^{(N+1)/2}} \Gamma\left(\frac{N+1}{2}\right) \zeta(N-1) T^{N-1}, \quad (18)$$

where $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ is the usual Riemann zeta function. Using the fact that the critical temperature of the condensate is reached when $\mu = M$, in $N = 3$ space dimensions, we have:

$$T_c = \left(\frac{3\rho}{M}\right)^{1/2}, \quad (19)$$

which coincides with the ultrarelativistic BEC results known in the literature [7-10].

Now, let us study the situation for the nonrelativistic limit which here means to take the low temperature limit, $\beta M \gg 1$. Using the asymptotic expansion for the Bessel function

$$K_\nu(x) \simeq \sqrt{\frac{\pi}{2x}} e^{-x}, \quad (20)$$

valid when $|x| \gg 1$ and $-\pi/2 < \arg x < \pi/2$, and taking $\beta M \gg 1$ in the free energy(8), we have ($\theta = 2n\pi$, $\sigma(\theta) = -1$):

$$\Omega(\beta, \mu) = -\frac{2V}{\beta} \left(\frac{M}{2\pi\beta}\right)^{\frac{N}{2}} \sum_{n=1}^{+\infty} \cosh(n\beta\mu) \left(\frac{1}{n}\right)^{1+\frac{N}{2}} e^{-n\beta M}. \quad (21)$$

In this regime the charge density reads

$$\rho(\beta, \mu) = 2 \left(\frac{M}{2\pi\beta}\right)^{\frac{N}{2}} \sum_{n=1}^{+\infty} \sinh(n\beta\mu) \left(\frac{1}{n}\right)^{\frac{N}{2}} e^{-n\beta M}. \quad (22)$$

Exactly as in the ultrarelativistic case, here the critical point is reached when $\mu \rightarrow M$. So, in the nonrelativistic limit, the condensation condition implies that $\beta\mu \gg 1$ and then the charge density reduces to

$$\rho(\beta, \mu) = \left(\frac{M}{2\pi\beta}\right)^{\frac{N}{2}} \sum_{n=1}^{+\infty} \left(\frac{1}{n}\right)^{\frac{N}{2}} e^{n\beta(\mu-M)}. \quad (23)$$

At the critical point $\mu = M$ we get simply

$$\rho = \zeta\left(\frac{N}{2}\right) \left(\frac{T_c M}{2\pi}\right)^{\frac{N}{2}}, \quad (24)$$

which for $N = 3$ space dimensions leads to the critical temperature

$$T_c = \frac{2\pi}{M} \left(\frac{\rho}{\zeta(3/2)}\right)^{2/3}, \quad (25)$$

in agreement with the nonrelativistic BEC results known in the literature [5-10].

Looking at the Eqs. (18) and (24) for the charge density in N space dimensions for the ultrarelativistic and nonrelativistic cases one can see that the condensate is not defined in two space dimensions for both cases, since the Riemann zeta function $\zeta(s)$ has a pole at $s = 1$. Hence, we find that the condensation for free massive fields (both in the ultra or nonrelativistic limits) occurs for space dimensions $N > 2$, which is in accordance with the BEC results for the massive case in the absence of external fields [5-10].

We presented here a relativistic partition function from which we described both the nonrelativistic and ultrarelativistic condensation. With our partition function we were also able to describe fermions and bosons at the same time by varying the parameters θ and σ . From this partition function we found the general condensation condition of the fields, *i.e.*, they should strictly obey *periodic* boundary conditions at finite temperature. This condition can be achieved by any field including fermions but conditioned to the fact that they must be coupled to topological fields which result in a total phase $\Theta = 2n\pi$. On the other side, bosons coupled to those fields in such a way that $\Theta \neq 2n\pi$ do not condense.

We also described here a kind of finite temperature bosonization (or fermionization) by including the arbitrary θ in the partition function. We have shown that no condensation occurs for fields with $\theta \neq 2n\pi$. However, a fermionic field interacting with a topological field $A_0 = \pi/\beta$ will also condense, so this should correspond to a bosonization process. Reversely, adding such a field to a boson we fermionize it once they will exclude rather than condense. Note that this picture is valid only at finite temperatures since at zero temperature the topological field $A_0 = \theta/\beta$ vanishes.

Our results also suggest that pairs of fermions can also condense as they would satisfy the condition $\Theta = 2n\pi$, as well as arbitrary combinations of θ -fields as long as that

condition is preserved. However, to have a clear picture of such situations one needs to consider some kind of interaction between these fields which was not considered in our approach. This is presently under investigation and will be reported elsewhere.

Acknowledgments. H.B.-F. acknowledges R. Jackiw for reading a preliminary version of the manuscript and for useful suggestions on it, and the hospitality of the Center for Theoretical Physics - MIT, where part of this work was done. The authors H.B.-F. and C.F. were partially supported by CNPq (Brazilian agency).

References

- [1] A. Einstein, S. B. Preus. Akad. Wiss. **22**, 261 (1924).
- [2] S. N. Bose, Z. Phys. **26**, 178 (1924).
- [3] F. London, Phys. Rev. **54**, 947 (1938).
- [4] M. H. Anderson, J. R. Ensher, M. R. Mathews, C. E. Wieman and E. A. Cornell, Science **269**, 198 (1995).
- [5] See, *e. g.*, K. Huang, *Statistical Mechanics*, Wiley, New York, 1963.
- [6] R. Bechmann, F. Karsch and D. E. Miller, Phys. Rev. Lett. **43**, 1277 (1979); C. A. Aragão de Carvalho and S. Goulart Rosa Jr., J. Phys. A **13**, 3233 (1980).
- [7] H. E. Haber and H. A. Weldon, Phys. Rev. Lett. **46**, 1497 (1981); Phys. Rev. D **25**, 502 (1982); J. Math. Phys. **23**, 1852 (1982).
- [8] J. I. Kapusta, Phys. Rev. D **24**, 426 (1981); for a review see J. I. Kapusta, *Finite temperature field theory*, Cambridge, 1989.
- [9] D. J. Toms, Phys. Rev. Lett. **69**, 1152 (1992); Phys. Rev. D **47**, 2483 (1993); *ibid.* **51**, 1886 (1995); Phys. Lett. B **343**, 259 (1995).
- [10] K. Kirsten and D. J. Toms, Phys. Rev. D **51**, 6886 (1995); Phys. Lett. B **368**, 119 (1996); Phys. Rev. D **55**, 7797 (1997).

- [11] S. Deser, R. Jackiw and S. Templeton, Phys. Rev. Lett. **48** (1982) 975; Annals of Phys. **140** (1982) 372.
- [12] For a review see F. Wilczek (ed.), *Fractional Statistics and Anyon Superconductivity*, World Scientific, Singapore, 1990.
- [13] P. Borges, H. Boschi-Filho and C. Farina, Mod. Phys. Lett. A **13**, 843 (1998).
- [14] D. P. Arovas, R. Schrieffer, F. Wilczek and A. Zee, Nucl. Phys. B **251** (1985) 117.
- [15] C. Bernard, Phys. Rev. D **9**, 3312 (1974).
- [16] A. Actor, Phys. Rev. D **27**, 2548 (1983); Ann. of Phys. (N.Y.) **159**, 445 (1985); Phys. Lett. B **157**, 53 (1985); Nucl. Phys. B **256**, 689 (1986).
- [17] T. Matsubara, Prog. Theor. Phys. **14**, 351 (1955).
- [18] H. Boschi-Filho, C. Farina and A. de Souza Dutra, J. Phys. A **28**, L7 (1995).
- [19] H. Boschi-Filho and C. Farina, Phys. Lett. A **205**, 255 (1995).
- [20] H.-T. Elze, W. Greiner and J. Rafelski, J. Phys. A **6**, L149 (1980).
- [21] P. Borges, H. Boschi-Filho and C. Farina, hep-th/9811166.